

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

International Edition

GDCh

www.angewandte.org

2009–48/44



## The key role of the interfaces ...


... in the electrode–molecule contact can be probed by studying the time evolution of molecular junctions. In their Communication on page 8273 ff., A. Filoramo and co-workers show that the “blinking” of the thiol–gold bond can be suppressed by an appropriate choice of molecule: devices that contain the terphenylene molecule exhibit a stochastic variation in the conductance, while devices that contain the terthiophene molecule are extremely stable over a period of several weeks (Illustration: D. Nozaki).

 WILEY-VCH

## Inside Cover

**Diana Dulić, Florian Pump, Stephane Campidelli, Pascal Lavie, Gianarelio Cuniberti, and Arianna Filoramo\***

**The key role of the interfaces** in the electrode–molecule contact can be probed by studying the time evolution of molecular junctions. In their Communication on page 8273 ff., A. Filoramo and co-workers show that the “blinking” of the thiol–gold bond can be suppressed by an appropriate choice of molecule: devices that contain the terphenylene molecule exhibit a stochastic variation in the conductance, while devices that contain the terthiophene molecule are extremely stable over a period of several weeks (Illustration: D. Nozaki).



## Controlled Stability of Molecular Junctions\*\*


Diana Dulić, Florian Pump, Stephane Campidelli, Pascal Lavie, Gianauelio Cuniberti, and Arianna Filoramo\*

Using single molecules as the building blocks of electronic devices is the ultimate goal of molecular electronics. However, measuring, understanding and manipulating the transport of charge through molecules attached to nanosized electrodes remain difficult tasks. It is thus important to fabricate molecular junctions exhibiting reproducible and stable electrical response and to evaluate their performance.<sup>[1,2]</sup> While chemists are able to synthesize molecules with amazing electronic, optical, magnetic, thermoelectric, or electromechanical properties and with the potential to provide electronic devices with novel functionality, integration into single-molecule devices remains the main problem of the field of “molecular electronics”. It has been demonstrated that electron transport through a molecule depends not only on its intrinsic properties but also on specific details of the contacts and the local environment.<sup>[3]</sup> Here we report a new strategy to monitor the conductance of such molecular junctions that enables further insight to be gained into the detailed nature of the conductance of single molecules. Due to the stability and reproducibility of our junctions, our method provides a new perspective on studies of electronic transport on the nanometer scale.

The first consistent results for the conductivity of a molecule in solution were obtained by statistical analysis of data collected by the scanning tunneling microscope (STM) break-junction method.<sup>[4]</sup> By repeatedly moving the STM tip into and out of contact with a molecular monolayer deposited on a substrate electrode, thousands of molecular junctions can quickly be created and statistical analysis permits the

conductance of the molecules with the most probable contact geometries to be determined. Another elegant method, the mechanically controllable break-junction (MCBJ) technique, has also been proposed to control the spacing between two metallic electrodes with subatomic ( $< 10^{-10}$  m) resolution.<sup>[5]</sup> However, the pioneering MCBJ experiments<sup>[6-9]</sup> did not employ statistical approaches, and therefore the number of molecules measured was uncertain.<sup>[4,7]</sup> Recently, first MCBJ experiments using the statistical approach on alkane dithiol molecules in solution were reported<sup>[10]</sup> and showed excellent agreement with earlier STM break-junction data.<sup>[11]</sup>

Here we study and analyze statistically electron transport in terthiophene (T3) and terphenylene (P3) molecules covalently bonded (via thiol groups) to two gold electrodes by the MCBJ technique in vacuum (Figure 1). To obtain the conductance for the most probable contact geometry, we first adopted a statistical approach similar to one described in the literature.<sup>[10]</sup> Details of the experiment are given in the Supporting Information. Compounds P3 and T3 were synthesized according to methods described in the literature.<sup>[12,13]</sup> Their structures are shown in Figure 1d and e, respectively. They were chosen because of their similar lengths, conjugation paths, and coupling groups, while the presence of additional sulfur atoms in the backbone of T3 modifies the orbital distribution and thus influences the transport characteristics.



**Figure 1.** a) SEM image of a break junction. b) Layout of the MCBJ technique. c) Schematic vision of the molecular arrangements in the break junction. d) and e) Conductance histograms (repetitional statistics) at  $V = 20$  mV, constructed from 100 traces for two different samples with P3 molecules and two different samples with T3 molecules, respectively; insets show typical conductance traces for P3 and T3.

[\*] Dr. D. Dulić,<sup>[†]</sup> Dr. S. Campidelli, P. Lavie, Dr. A. Filoramo  
CEA, Laboratoire d'Electronique Moléculaire, IRAMIS/SPEC  
91191 Gif sur Yvette cedex (France)  
Fax: (+33) 1-6908-6640  
E-mail: arianna.filoramo@cea.fr  
Homepage: <http://iramis.cea.fr/spec/lem>  
F. Pump, Prof. Dr. D. G. Cuniberti  
Institute for Materials Science and Max Bergmann Center of  
Biomaterials, Dresden University of Technology, 01062 Dresden,  
(Germany)

[†] Present address: Institute of Physics  
Pregrevica 118, 11080 Belgrade (Serbia)

[\*\*] We thank G. Blake and R. Ferreira for useful suggestions and careful reading of this text. This project is partially funded by Marie Curie Intra-European Fellowship EIF N° 025839, by the VolkswagenStiftung, and by the DFG Priority Programme 1243 “Quantum transport at the molecular scale. G.C. and F.P. acknowledge the ZIH at the Dresden University of Technology for computational resources.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200902168>.


For each sample we performed a series of about 100 opening/closing cycles while monitoring the current (for details see Supporting Information). From now on we refer to this method as repetitional statistics, since each of the individual files used to build the histogram represents a molecular junction with a different conformation with respect to the electrodes. Representative closing traces for P3 and T3 are shown in the insets of Figure 1 d and e, respectively. The observed conductance plateaus are remarkably long, and their lengths are roughly those of the molecules (ca. 1.4 nm for P3 and 1.2 nm for T3).

In the case of T3 (Figure 1 e), the conductance histograms of different MCBJ samples have similar peak values but different peak widths (single Gaussian fit). The P3 histograms (Figure 1 d) contain more structure than those of T3. However, as reported in the literature, this robust statistical method provides a reliable value for the conductance but at the expense (because of peak widths) of fine structure in the main peak, which gives information about fluctuations in molecular junctions.<sup>[14–16]</sup>

Since our MCBJ electrodes can be controlled at subatomic resolution and has impressive stability in the presence of the molecules (on the order of weeks), we developed a new statistical approach referred to hereafter as fluctuational statistics, which enables the time evolution of molecular junctions to be monitored and thus fully complements the repetitional statistics data. As shown below, this method allows the observed conductance peaks corresponding to the fine structure of the main peak to be mapped. To implement it we blocked the interelectrode distance at the position of a conductance plateau by stopping the motor and then measured the current through the junction as a function of time (left panels of Figure 2; for details, see Supporting Information).

There is one major difference between the P3 and T3 molecular junctions. The conductance of the molecule changes, typically every few minutes, between two distinct levels for P3 (Figure 2a) but not for T3 (Figure 2b). The corresponding conductance histograms are traced in the right panels of Figure 2. Stochastic variations in conductance have previously been observed in thiol-terminated molecules by STM or the conductive atomic force microscopy imaging technique.<sup>[17–20]</sup> They were related to “blinking” of the thiol-gold bond, an intrinsic feature of any molecule connected to a gold surface via thiol groups.<sup>[18]</sup> However, the T3 molecule does not show such well-defined “quantized changes” in conductance.

The conductance switching observed for P3 may be due to a thermally induced change in the bonding sites of the thiol groups terminating the molecules on the electrode surfaces. It is known that the sulfur atoms prefer to bind at high-symmetry positions on the gold surface (top, bridge, hollow fcc, and hollow hcp sites, assuming a (111) surface), and that the bonding site has an important influence on the conduction properties of single molecules between metallic electrodes.<sup>[21,22]</sup> Thus, we computed the transmission function through the P3 molecule attached to gold electrodes using the gDFTB method,<sup>[23]</sup> a combination of the non-equilibrium



**Figure 2.** Left panels: Conductance versus time for P3 (a) and T3 (b) molecules. The circled regions are zoomed in the insets. Data points recorded every 0.5 s at 100 mV bias voltage. Right panels: corresponding fluctuational histograms (violet for P3 and green for T3). They are compared with the repetitional histograms in the insets (pink for P3 and orange for T3).

Green function technique and the density-functional-based tight-binding method DFTB.<sup>[24,25]</sup>

Comparing the zero-bias conductance values (defined as  $G = G_0 T(E_F)$  with  $G_0 = 2e^2 h$ ; for more details and the corresponding transmission curves, see Supporting Information) obtained for the four different bonding sites, we find that the bonding site has a huge influence on the conductance (Table 1).


**Table 1:** Calculated conductance of P3 molecules for different bonding sites between sulfur atoms and Au(111) surface.

Bonding site	$G [10^{-2} G_0]$	Bonding site	$G [10^{-2} G_0]$
hollow fcc	0.18	top	6.16
hollow hcp	0.19	bridge	0.32

It is likely that the thermal motion of the connecting sulfur atom on the electrode surface at room temperature can vary the conductance through the molecule, and the stability of the bonding at the high-symmetry points could be the reason for the distinct peaks identified in the statistical analysis in Figure 2a. However, for T3 the presence of a sulfur atom in the ring could increase the overall stability of the bonding<sup>[26]</sup> and thus explain the absence of observable “blinking” in the

junction. Note that the fluctuational histograms show improved resolution compared to the repetitional ones, as shown in the right insets of Figure 2.

As shown in Figure 3, the analysis of the conductance on longer timescales and at different voltages reveals further interesting features. The P3 molecule again exhibits a richer



**Figure 3.** a) and b) Conductance versus time (insets) with changing bias voltage and comparison (main figures) between repetitional (pink and orange) and fluctuational (violet and green) statistics for P3 and T3 respectively. Points were taken every 0.5 s. For P3, stochastic switching is observed between 110 and 190 h.

evolution of conductance (inset of Figure 3a). In particular, between 110 and 190 h the conductance oscillates between two distinct levels (independent of the applied voltage). The fact that this switching behavior does not appear immediately after formation of the junction suggests a local voltage/current-activated effect.<sup>[27]</sup> More generally, all molecular junctions drift over long timescales due to progressive and gradual rearrangement of the molecules and charge distribution. As discussed below, the comparison between repetitional and fluctuational statistics sheds light on the conductance evolution, as the local environment of the connected molecule(s) is modified.

The fluctuational histogram for P3 exhibits distinct peaks at 0.15, 0.20, 0.25, and 0.46  $G_0$  (violet histogram in Figure 3a). These values are consistent with those we obtained from the repetitional statistics (pink histogram in Figure 3a) and correspond to either hopping between hollow-bridge configurations or additional trapping of molecule(s). Differences between the two statistical approaches are more evident on longer timescales. In particular, the field-induced “wearing on” of the junction by surrounding molecules becomes more important. We emphasize that the time evolution of T3 conductance does not exhibit any switching, even though different bonding-site configurations were eventually

observed in the repetitional approach.<sup>[28]</sup> As reported in Figure 3b (inset), on varying the voltage the conductance versus time changed from 0.1  $G_0$  to 0.2  $G_0$  before increasing further, never returning to the initial value. We interpret this observation by postulating that near-integer multiples of the fundamental conductance correspond to the formation of a multiple molecular junction in which a small number of molecules are connected in parallel between the electrodes, that is, two molecules are bridging the junction.<sup>[4]</sup> In these data, note an abrupt increase in conductance to 0.2  $G_0$  at 200 mV bias voltage. To confirm that increasing the voltage promotes collection of another molecule we formed a fresh molecular junction and repeated the experiment at 100 mV (see Supporting Information).

Our conductance values are substantially higher than those obtained by the repetitional statistics STM break-junction method in the liquid phase for similar molecules.<sup>[14]</sup> Theoretical calculations<sup>[22]</sup> show that the conductivity of a molecular junction is extremely sensitive to the geometry of the contact, which in our case is different from the STM geometry, and the presence of different media might play a role. Moreover, we do not claim that the most probable conductance value obtained by the repetitional statistics method corresponds solely to the conductance of a single molecule or exclude the possibility that the overall conductance is influenced by the presence of neighboring molecules. However, our data show that the junction persists for a long time only when the bridging molecule(s) is (are) connected to both electrodes, and that the conductance obtained by fluctuational statistics corresponds to the most probable value obtained by the repetitional statistics method. Finally, note that the stochastic changes in STM imaging experiments<sup>[19]</sup> do not occur in experiments with larger numbers of the same molecules.<sup>[29,30]</sup>

In conclusion, we have studied the influence of the interface on simple molecular wires, a factor essential to understand before molecular electronic devices can be realized. Due to the week-long stability and reproducibility of our junctions, our method opens new horizons in studying molecules with more interesting intrinsic properties, as well as designing and studying molecules with different end groups that could suppress stochastic switching. This will have an impact not only on the field of single-molecule electronics, but also in general molecular electronics dealing with self-assembled monolayers.

Received: April 22, 2009

Revised: July 20, 2009

Published online: September 2, 2009

**Keywords:** break-junction method · electron transport · molecular electronics · scanning probe microscopy · single-molecule studies

[1] N. J. Tao, *Nat. Nanotechnol.* **2006**, *1*, 173–181.

[2] *Introducing Molecular Electronics* (Eds G. Cuniberti, G. Fagas, K. Richter), Springer, Berlin, **2005**.

[3] M. del Valle, R. Gutierrez, C. Tejedor, G. Cuniberti, *Nat. Nanotechnol.* **2007**, *2*, 176–179.

- [4] B. Q. Xu, N. J. Tao, *Science* **2003**, *301*, 1221–1223.
- [5] J. M. van Ruitenbeek, A. Alvarez, I. Pineyro, C. Grahmann, P. Joyez, M. H. Devoret, D. Estève, C. Urbina, *Rev. Sci. Instrum.* **1996**, *67*, 108–111.
- [6] M. A. Reed, C. Zhou, C. J. Muller, T. B. Burgin, J. M. Tour, *Science* **1997**, *278*, 252–254.
- [7] C. Kergueris, J.-P. Bourgoin, S. Palacin, D. Esteve, C. Urbina, M. Magoga, C. Joachim, *Phys. Rev. B* **1999**, *59*, 12505–12513.
- [8] J. Reichert, R. Ochs, D. Beckmann, H. B. Weber, M. Mayor, H. von Löhneysen, *Phys. Rev. Lett.* **2002**, *88*, 176804.
- [9] D. Dulić, S. J. van der Molen, T. Kudernac, H. T. Jonkman, J. J. D. de Jong, T. N. Bowden, J. van Esch, B. L. Feringa, B. J. van Wees, *Phys. Rev. Lett.* **2003**, *91*, 207402.
- [10] M. T. González, S. Wu, R. Huber, S. J. van der Molen, C. Schönenberger, M. Calame, *Nano Lett.* **2006**, *6*, 2238–2242.
- [11] X. Li, J. He, J. Hihath, B. Xu, S. M. Lindsay, N. Tao, *J. Am. Chem. Soc.* **2006**, *128*, 2135–2141.
- [12] B. de Boer, H. Meng, D. F. Perepichka, J. Zheng, M. M. Frank, Y. J. Chabal, Z. Bao, *Langmuir* **2003**, *19*, 4272–4284.
- [13] D. A. Krapchetov, H. Ma, A. K. Y. Jen, D. A. Fischer, Y.-L. Loo, *Langmuir* **2005**, *21*, 5887–5893.
- [14] L. Venkataraman, J. E. Klare, C. Nuckolls, M. S. Hybertsen, M. L. Steigerwald, *Nature* **2006**, *442*, 904–907.
- [15] R. Huber, et al., *J. Am. Chem. Soc.* **2008**, *130*, 1080–1084.
- [16] J. Hihath, N. Tao, *Nanotechnology* **2008**, *19*, 265204.
- [17] P. A. Lewis, C. E. Inman, Y. Yao, J. M. Tour, J. E. Hutchison, P. S. Weiss, *J. Am. Chem. Soc.* **2004**, *126*, 12214–12215.
- [18] W. A. Wassel, R. R. Fuijrer, N. Kim, C. B. Gorman, *Nano Lett.* **2003**, *3*, 1617–1620.
- [19] G. K. Ramachandran, T. J. Hopson, A. M. Rawlett, L. A. Nagahara, A. Primak, S. M. Lindsay, *Science* **2003**, *300*, 1413–1416.
- [20] I.-W. P. Chen, M.-D. Fu, W.-H. Tseng, J.-Y. Yu, S.-H. Wu, C.-J. Ku, C.-H. Chen, S.-M. Peng, *Angew. Chem.* **2006**, *118*, 5946–5950; *Angew. Chem. Int. Ed.* **2006**, *45*, 5814–5818.
- [21] A. Grigoriev, J. Sköldbberg, G. Wending, Z. Crljen, *Phys. Rev. B* **2006**, *74*, 045401.
- [22] H. Basch, R. Cohen, M. A. Ratner, *Nano Lett.* **2005**, *5*, 1668–1675.
- [23] A. Di Carlo, A. Pecchia, L. Latessa, T. Frauenheim, G. Seifert in *Introducing Molecular Electronics* (Eds.: G. Cuniberti, G. Fagas, K. Richter), Springer, Berlin, **2005**, pp. 153–184.
- [24] M. Elstner, D. Porezag, G. Jungnickel, J. Elsner, M. Haugk, T. Frauenheim, S. Suhai, G. Seifert, *Phys. Rev. B* **1998**, *58*, 7260–7268.
- [25] T. Frauenheim, G. Seifert, M. Elstner, Z. Hajnal, G. Jungnickel, D. Porezag, S. Suhai, R. Scholz, *Phys. Status Solidi B* **2000**, *217*, 41–62.
- [26] B. de Boer, H. Meng, D. F. Perepichka, J. Zheng, M. M. Frank, Y. J. Chabal, Z. Bao, *Langmuir* **2003**, *19*, 4272–4284.
- [27] Z. Huang, B. Xu, Y. Chen, M. Di Ventra, N. Tao, *Nano Lett.* **2006**, *6*, 1240–1244.
- [28] The presence of the sulfur atom in the thiophene ring stabilizes the linkage with the electrodes and avoids thermally activated fluctuations among different bonding-site configurations. In the repetitional approach, the possibility of observing different bonding sites is not affected in the same way since a new junction is formed for each opening/closing cycle.
- [29] L. Cai, M. A. Cabassi, H. Yoon, O. M. Cabarcos, C. L. McGuinness, A. K. Flatt, D. L. Allara, J. M. Tour, T. S. Mayer, *Nano Lett.* **2005**, *5*, 2365–2372.
- [30] J. G. Kushmerick, D. B. Holt, J. C. Yang, J. Naciri, M. H. Moore, R. Shashidhar, *Phys. Rev. Lett.* **2002**, *89*, 086802.